

LOGAN

LIQUID NITROGEN - FERTILIZER ORGANIC

ANALYSIS	w/w	g/L
Organic nitrogen (N)	6%	76.2
Organic carbon (C) of biological origin	25.2%	320

ALLOWED IN ORGANIC FARMING IN ACCORDANCE WITH THE EUROPEAN REGULATION

DOSAGE AND INSTRUCTIONS FOR USE

Foliar application:

- **Fruit trees (pome and stone fruits) and table grape:** during the vegetative stage to improve plant growth, 3-4 applications every 7-10 days 5-2.5 L/ha
- **Table grape:** one application with bunches 3-4 cm and repeat after 6-7 days with the aim of promoting bunch elongation 1.5 L/ha
- **Tropical fruit trees (mango, banana, papaya, etc.) and citrus:** during new leaf flushing and at flowering, 2-3 applications every 7-10 days 1.5-2.5 L/ha
- **Vegetable and industrial crops (potato, sunflower, sugar cane, cotton, etc.):** during the whole cycle, 3-4 applications every 7-10 days 1.5-2 L/ha
- **Cereals (rice, wheat, corn, etc.):** 1st application at post-emergence usually combined with herbicides; 2nd application at pre-flowering stage 1-2 L/ha
- **Flower and ornamental crops:** during the whole crop cycle, 2-3 applications every 7-10 days 100-200 mL/hL

Dosages are calculated for normal water distribution volumes

Handling and storage



Security and ecological information

In case of spillage, collect with sawdust and/or sand
Keep out of reach of children



warning

Keep away from flames, and store in a cool, well-ventilated place. Avoid prolonged exposure to low temperatures. Check container periodically. In case of accidental spill contain the leak, avoid water contamination and, whenever necessary, dispose according to the laws in force. **SHAKE BEFORE USE**

PACKING

	1 PALLET CANS 25 LT 48 cans= 1200 Lt			
CAN 25 LT			IBC 1000 lt	